

IMSERSO / European Centre / UNECE

# Workshop

---

## **Sustainable Ageing Societies: Indicators for Effective Policy-Making**

### **Thematic session 1a**

Demographic challenges,  
population ageing, income and wealth

Madrid, Spain, 14-16 April 2004

## **Monitoring Morbidity, Mortality and Longevity Developments in the UNECE Region, and their Effect on Long Term Population Trends**

*Jean-Marie Robine*

## Sustainable ageing societies: Indicators for Effective Policy-Making

*IMSERMO Workshop, Madrid 14-16.04.2004*

### Policy brief: **Monitoring Morbidity, Mortality and Longevity Developments in the UNECE Region, and their effect on long term population trends: the most policy-relevant indicators**

By Jean-Marie Robine\* and Carol Jagger\*\*

\*Demographer, INSERM, University of Montpellier 1, robine@valdorel.fnclcc.fr

\*\*Epidemiologist, University of Leicester

The aim of this paper is to provide the essential guidelines for development of comparative social indicators in the domains of morbidity, mortality and longevity in the UNECE Region, and their effect on long-term population trends. The set of indicators should be relatively stable over time to ensure continuity. We propose guidelines that fulfil the main criteria listed for EU social indicators; that is


- each indicator should be policy relevant at the EU level;
- comparable across EU member states;
- available using harmonized sources;
- measurable over time;
- easily understood;

## 1 Demographic and epidemiological transition in the UNECE Region

### 1.1 An ageing population and the emergence of the chronic diseases

The fall in infant mortality during the 19<sup>th</sup> and early 20<sup>th</sup> century, a result of increased food availability, decline in infectious diseases and improvement of hygiene, gave a greater number of newborn the opportunity to live a "normal" lifespan (Kannisto, 2001). This epidemiological transition (Omran, 1971) led to population ageing and to the emergence of chronic morbidity. In this context, ageing-related diseases and their consequences became an important social issue, resulting in the *International Classification of Impairments, Disabilities, and Handicaps: A manual of classification relating to the consequences of disease* (ICIDH), published by WHO in 1980 (see Figure 1).

**Figure 1: International Classification of Impairments, Disabilities and Handicaps (ICIDH): Integration of concepts (WHO, 1980)**




The study of the disablement process during the 1980s and the 1990s resulted in several new concepts: functional limitation, activity restriction, unmet need, physical dependency, loss of

autonomy, social integration, and quality of life, and many indicators have been proposed to measure these concepts.

One of the most important questions was whether total life expectancy was increasing faster or slower than life expectancy in good health (i.e. without chronic disease, functional limitation, activity restriction). This question led to the development of a new family of indicators, health expectancies. The general model of health transition proposed by WHO in 1984 (see Figure 2) distinguished life expectancy, disability-free life expectancy and morbidity-free life expectancy, and its strength lies in its ability to assess the likelihood of different health scenarios: a pandemic of chronic diseases and disabilities (Gruenberg, 1977; Kramer, 1980), a compression of morbidity (Fries, 1980), contradictory evolutions including the scenario of dynamic equilibrium (Manton, 1982), or a postponement of all morbid events (diseases, disabilities and mortality) at older ages (Strehler, 1975).

**Figure 2: The general model of health transition (WHO, 1984). Observed mortality and hypothetical morbidity and disability survival curves for females, United States of America, 1980.**



The health expectancy approach has now been extended to incorporate the modern concepts of the disablement process: chronic morbidity, functional limitation, activity restriction, physical dependency and progressively more attention has been devoted to mental health expectancy (Robine and Jagger, 2003).

## 1.2 The lengthening of life and the emergence of extremely old persons

In addition to the reduction in infant mortality, social and economic developments contributed to a reduction in the mortality of older people, beginning shortly after WWII in low mortality countries. Generally this was a result of the considerable improvement in living conditions experienced by these cohorts after WWII, including the wider availability of health and medical services. When this phase began under the impact of economic, medical and technical progress, it initially applied to a population in bad health in comparison with the current population of the same age. Thus the fall in mortality increased the prevalence of chronic diseases and disability by first increasing the survival rates of sick persons. This is the morbidity expansion scenario proposed by Gruenberg (1977) and Kramer (1980) which may explain the picture in the United States of America in the 1960s or 1970s. Gradually, medical progress slowed down the progression of chronic diseases involving a kind of equilibrium with the mortality decline, as proposed by Manton in 1982, for example the cardiovascular revolution (Meslé and Vallin, 2000). Progressively, the health of older people improved with the arrival of new cohorts who have been exposed longer to the social, economic, medical and technical advances resulting in a compression of morbidity, a phase which Austria, France or

the United States are currently experiencing today. However continued progress, in particular in the built environment and services provided to the oldest-old along with further fall in mortality has led to the emergence of extremely old populations such as super-centenarians and to new concepts like frailty (Fried et al, 2001). These new phenomena could result in a return to an expansion of morbidity (Robine and Michel, 2004).

In summary the UNECE Region is characterized by four elements:

- an increase in the survival rates of sick persons;
- a control of the progression of chronic diseases;
- an improvement in the health status and health behaviours of new cohorts of old people;
- an emergence of very old and frail populations.

All these elements coexist today and future scenarios will depend on their respective weights. All should be monitored.

## **2. Essential guidelines for the development of indicators for monitoring morbidity, mortality and longevity developments in the UNECE Region**

There are three main points that require careful examination when proposing such indicators. They are (i) the domains to be covered, (ii) the technical characteristics of the indicators, and (iii) the data collection process.

### **2.1 Domains to be covered**

The broad domains to be covered have already been summarised above and are the demographic and epidemiologic history of the UNECE Region. In more detail they are:

- Longevity: referring to life duration;
- Chronic morbidity and co-morbidity: referring to diseases and ultimately causes of death. Co-morbidity is a key issue for the oldest-old;
- Disability: Referring to functional limitations and activity restrictions. The distinction between functioning at the body level and activity at an individual level is essential to properly monitor the disablement process associated with ageing and lengthening of life and it provides a framework for health interventions aiming to prevent, limit or slow down the disablement process (WHO, 1999). It is also implicated in;
- Independent life, autonomy, social integration: underlining the three main dimensions of activity at the society or community level (i) physical independence in daily activities (personal care, household care, work and leisure), (ii) mental autonomy in daily decisions and (iii) social integration;
- Frailty: Referring to the ability to mobilize resources (physical and mental) to cope with stress. It is recognized as a distinct entity from comorbidity and disability (Fried et al. 2004)

Loss of resources (weight, muscle, memory, vision) are the main characteristics of the current oldest-old;

- Quality of life: The main target is to increase the quality of life not merely the quantity.

## 2.2 Technical characteristics of the indicators

Having broadly defined the main domains, there are several technical characteristics of note:

- Prevalence, ratios and numbers: For specific domains, it is important to have not only ratios, such as the prevalence of disability per 100 or 1000 persons but also the crude numbers. With the increase in life expectancy, some numbers, for instance the number of nonagenarians, centenarians, frail or demented persons, are increasing rapidly and are key for those planning health and social care services. In addition speed indicators, such as the Centenarian Doubling Time (CDT) are helpful. Special attention should be paid to the choice of denominators, whether current population or cohort of origin;

- Measures of central tendency and dispersion: Not all older people are sick, frail or disabled and as well as the average (mean or modal) level of disability, it is also important to know the distribution of the population through the various continuums. Specifically with regard to longevity, the mode of life durations, indicating the most typical life durations, appear to be much more relevant than life expectancy at birth which confounds premature mortality with longevity. Life expectancy at birth is a poor indicator of the most typical life durations;

- Health expectancies: We need simple global indicators, easy to understand, combining mortality and morbidity, longevity and quality of life to answer the basic questions raised by the demographic and epidemiological transition in the UNECE Region (see above). Such a set of indicators has been proposed by the Euro-REVES group for the EU Health Monitoring Program (Robine and Jagger, 2003) and will be regularly computed within the new EU Public Health Programme (2003-2008).

## 2.3 Data collection

Finally, to accurately monitor morbidity, mortality and longevity through a set of indicators, there are some essential issues in the data collection:

- Harmonization of the survey instruments: Monitoring morbidity, mortality and longevity developments in the UNECE Region implies harmonised national health and ageing surveys. Current initiatives such as SHARE (Survey of Health, Ageing and Retirement in Europe), partially comparable with ELSA (English Longitudinal Survey on Ageing) and HRS (Health and Retirement Survey in the United States) and the preparation of EHIS (European Health Interview Survey) by Eurostat are moving forwards to this end;

- Health examination component including cognitive physical tests: The introduction of a health examination component (with physical and cognitive tests) in health and ageing surveys will increase comparability. This may also be achieved through the development of other methods such as vignettes that allow interviewees to rank the same “clinical” cases. All such techniques will increase bridge information for post-harmonisation and calibration to complement pre-harmonisation;

- Cross-sectional surveys versus panel studies: During the last decade, several panel studies on ageing allowed monitoring of the incidence of health problems and computed more

sophisticated period health expectancies using a multi-state life table approach. However basic monitoring requires first repeating cross-sectional surveys to ensure the continuity of chronological series. Panel studies are not without methodological problems including attrition, selection and wave effects such as learning effects for cognitive tests.

## **Towards an UNECE Ageing Survey**

The development of a regular UNECE ageing survey following these guidelines would contribute most of the social indicators needed to monitor morbidity, mortality and longevity developments in the UNECE Region.

## **References**

- Fried LP, Tangen CM, Walston J, et al (2001) Frailty in older adults: evidence for a phenotype. *J Gerontol* 56A(3):M146-56.
- Fried LP, Ferrucci L, Darer J, Williamson JD, Anderson G (2004) Untangling the concepts of disability, frailty and comorbidity: implications for improving targeting and care. *J Gerontol* 59A(3):M255-63.
- Fries JF (1980) Aging, natural death, and the compression of morbidity. *N Engl J Med* 303:130-135.
- Gruenberg EM (1977) The failures of success. *Milbank Memorial Fund Quarterly/Health Society* 55:3-24.
- Kanisto V (2001) Mode and dispersion of the length of life. *Population* 13(1):159-71.
- Kramer M (1980) The rising pandemic of mental disorders and associated chronic diseases and disabilities. *Acta Psychiatrica Scandinavica* 62 (Suppl. 285):282-297.
- Manton KG (1982) Changing concepts of morbidity and mortality in the elderly population. *Milbank Memorial Fund Q / Health Soc* 60:183-244.
- Meslé F, Vallin J (2000) Transition sanitaire: tendances et perspectives. *Médecine/Sciences* 16:1161-71.
- Omran AR (1971) The epidemiological transition: a theory of epidemiology of population change. *Milbank Mem Fund Q* 49:509-38.
- Robine JM, Jagger C, the Euro-REVES group (2003) Creating a coherent set of indicators to monitor health across Europe: the Euro-REVES 2 project. *European Journal of Public Health*, 13(3):6-14.
- Robine JM, Michel JP (2004) Looking forward to a general theory on population aging, *J Gerontol* 59A(x): Mxxx-xxx in press.
- Strehler BL (1975) Implications of aging research for society. In: Theoretical concepts of developmental and age changes. *Federation Proceedings* 34(1):5-8.
- World Health Organization (1980) *International Classification of Impairments, Disabilities, and Handicaps: A manual of classification relating to the consequences of disease*. WHO: Geneva.
- World Health Organization (1984) *The uses of epidemiology in the study of the elderly: Report of a WHO Scientific Group on the Epidemiology of Aging*. Geneva: WHO (Technical Report Series 706).
- World Health Organization (1999) *International Classification of Functioning and Disability (ICIDH 2-Beta 2)*. Geneva: WHO.

## **Key words**

EPIDEMIOLOGICAL TRANSITION / POPULATION AGEING / CHRONIC MORBIDITY / DISABLEMENT PROCESS / FUNCTIONAL LIMITATION / ACTIVITY RESTRICTION / UNMET NEED / PHYSICAL DEPENDENCY / LOST OF AUTONOMY / SOCIAL INTEGRATION / QUALITY OF LIFE / HEALTH EXPECTANCY / DISABILITY-FREE LIFE EXPECTANCY / MORBIDITY-FREE LIFE EXPECTANCY / MENTAL HEALTH / LENGTHENING OF LIFE / PANDEMIC OF CHRONIC DISEASES AND DISABILITIES / COMPRESSION OF MORBIDITY / DYNAMIC EQUILIBRIUM / CARDIOVASCULAR REVOLUTION / BUILT ENVIRONMENT / SERVICES FOR THE OLDEST-OLD / EMERGENCE OF EXTREMELY OLD POPULATIONS / NONAGENARIANS / CENTENARIANS / SUPERCENTENARIANS / EMERGENCE OF FRAIL POPULATIONS / FRAILITY

